Perihelion precession

Perihelion precession

$$\alpha = \frac{6\pi GM}{c^2 a(1 - e^2)} \qquad \qquad \text{per single orbital period}$$

Perihelion precession

$$\alpha = \frac{6\pi GM}{c^2 a(1 - e^2)} \qquad \text{per single orbital period}$$
$$\Omega_{\text{GR}} = \frac{\alpha}{T} = \frac{3c}{a(1 - e^2)} \left(\frac{GM}{c^2 a}\right)^{3/2} \qquad \text{angular frequency}$$

Perihelion precession

$$\alpha = \frac{6\pi GM}{c^2 a(1 - e^2)} \qquad \text{per single orbital period}$$
$$\Omega_{\text{GR}} = \frac{\alpha}{T} = \frac{3c}{a(1 - e^2)} \left(\frac{GM}{c^2 a}\right)^{3/2} \qquad \text{angular frequency}$$

Mercury

 $\Omega_{\rm GR} = 43'' \cdot (100 \, \rm yr)^{-1}$

$$\Omega_{\text{planets}} \approx 500'' \cdot (100 \,\text{yr})^{-1}$$

Perihelion precession

$$\alpha = \frac{6\pi GM}{c^2 a(1 - e^2)} \qquad \text{per single orbital period}$$

$$\Omega_{\text{GR}} = \frac{\alpha}{T} = \frac{3c}{a(1 - e^2)} \left(\frac{GM}{c^2 a}\right)^{3/2} \qquad \text{angular frequency}$$

Mercury

PSR B1913+16

 $\Omega_{\rm GR} = 43'' \cdot (100 \, \rm yr)^{-1} \qquad \qquad \Omega_{\rm GR} \approx 4^\circ \cdot \, \rm yr^{-1}$

 $\Omega_{\text{planets}} \approx 500'' \cdot (100 \, \text{yr})^{-1}$

Perihelion precession

$$\alpha = \frac{6\pi GM}{c^2 a(1 - e^2)} \qquad \text{per single orbital period}$$
$$\Omega_{\text{GR}} = \frac{\alpha}{T} = \frac{3c}{a(1 - e^2)} \left(\frac{GM}{c^2 a}\right)^{3/2} \qquad \text{angular frequency}$$

MercuryPSR B1913+16S2 star in Sgt A*
$$\Omega_{GR} = 43'' \cdot (100 \text{ yr})^{-1}$$
 $\Omega_{GR} \approx 4^\circ \cdot \text{ yr}^{-1}$ $\Omega_{GR} \approx 12' \cdot (16 \text{ yr})^{-1}$

 $\Omega_{\text{planets}} \approx 500'' \cdot (100 \, \text{yr})^{-1}$

Schwarzschild in Kruszkal-Szekeres coordinates



Schwarzschild + interior solution in Kruszkal-Szekeres coordinates



Black holes

Black holes

regions of spacetime from which the communication with the observers at infinity is cut off due to extreme gravity. Typically containing singularities.

Black holes

regions of spacetime from which the communication with the observers at infinity is cut off due to extreme gravity. Typically containing singularities.

from far away they seem like ordinary massive objects, but without a surface

Black holes

regions of spacetime from which the communication with the observers at infinity is cut off due to extreme gravity. Typically containing singularities.

from far away they seem like ordinary massive objects, but without a surface

matter can fall into a BH (finite cross-section for absorption)

Black holes

regions of spacetime from which the communication with the observers at infinity is cut off due to extreme gravity. Typically containing singularities.

from far away they seem like ordinary massive objects, but without a surface

matter can fall into a BH (finite cross-section for absorption)

they are dark, but matter around BH may be hot and visible

Black holes

regions of spacetime from which the communication with the observers at infinity is cut off due to extreme gravity. Typically containing singularities.

from far away they seem like ordinary massive objects, but without a surface

matter can fall into a BH (finite cross-section for absorption)

they are dark, but matter around BH may be hot and visible

stellar mass black holes form from the collapse of the cores of massive stars

Black holes

regions of spacetime from which the communication with the observers at infinity is cut off due to extreme gravity. Typically containing singularities.

from far away they seem like ordinary massive objects, but without a surface

matter can fall into a BH (finite cross-section for absorption)

they are dark, but matter around BH may be hot and visible

stellar mass black holes form from the collapse of the cores of massive stars

Non-rotating BH described by the Schwarzschild solution, rotating BH by the Kerr solution

Introduction to general relativity

Mikołaj Korzyński (CFT PAN)

Winter semester 2023/24

Tuesdays, 9am-11am

This is the end of the course